

Association of Salivary Flow Rate & pH of Diabetes Mellitus Type II Subjects with Dental Caries and Gingivitis

Wasif Iqbal¹, Shafaq Noori², Abdur Rehman³

ABSTRACT

Previously it is suggested that diabetes mellitus may be the major cause of oral diseases. We aimed to assess the gingival and caries indices in type II Diabetes patients and also their correlation with salivary pH and flow rate. For this purpose we collected 100 subjects and divided into: Group I is control group contain 25 subjects; Group II is diabetes contain 75 subjects. Oral examination is asses after confirming the fasting blood sugar level. Mean fasting blood sugar level was 263 ± 62.7 in diabetes group. Mean gingival indices was 2.28 ± 0.62 and dental caries indices was 2.25 ± 0.61 in diabetes group. A significant differences were found between salivary flow rate and salivary pH with gingival and dental caries indices ($p < 0.05$). We found negative correlation of salivary pH of diabetes patients with gingival and caries indices, positive correlation of salivary flow rate of diabetes patients with gingival and caries indices and strong and positive correlation of caries and gingival indices. We concluded that low salivary PH in type II diabetes mellitus patients may cause gingivitis and dental caries and inflammation of gingival tissues may be secondary to caries.

Key Words: Gingivitis, dental caries, diabetes mellitus type II

(Citation: Iqbal Wasif ¹, Noori Shafaq ², Rehman Abdur ³
MedChannel 2026, 32 (4) 5-9

ARTICLE INFORMATION

Received on: 04-05-2021

Correction on: 11-12-2022

Accepted: 30-12-2022

Authors Contribution:

All authors contributed to the study. The lead author spearheaded study design, data analysis, and manuscript drafting. Authors two through five provided expertise in various aspects such as statistical analysis, clinical guidance, literature review, and data interpretation. The sixth author assisted in patient recruitment, data collection, and organization. The seventh author contributed to manuscript preparation and revision. All authors reviewed and approved the final manuscript for publication.

Conflict of Interest Disclosures: None

Disclose of Funding: None

Author Affiliation:

1. Associate Professor & Head Dental Material department, Muhammad Bin Qasim medical & dental college Karachi Pakistan.
2. Assistant Professor & Head Biochemistry department, Muhammad Bin Qasim medical & dental college Karachi Pakistan.
3. Dr. Abdur Rehman: Assistant Professor Dental Material department, Hamdard College of medicine & dentistry, Hamdard University Karachi.

Corresponding Author:

Dr. Hina Mudassir:

Assistant Professor Biochemistry
Department, Federal Urdu University of
Science & Technology Karachi.
Email: abdurrehmanmnds@gmail.com

INTRODUCTION

Diabetes mellitus is a common chronic disease that has emerged as a major health –care problem. People with diabetes are at an increased risk of developing oral conditions such as gingivitis, periodontal diseases and alveolar bone loss, which has been associated with persistent poor glycemic control. Periodontal disease can lead to recession of the gingival margin, which can expose more tooth surfaces to caries attack. People with diabetes can also experience hyposalivation and may suffer from salivary dysfunction. Oral hygiene is of prime importance to all individuals. Oral health priorities seek to reduce the negative impact of oral disease and their consequences.

Gingival inflammation is a common clinical feature detected in children and adults. It is characterized by swelling, redness and bleeding at the gums and it is described as an inflammatory reaction upon pro inflammatory cytokines that modulate the balance between humoral and cell associated immune responses. It is evident that plaque induced gingivitis is prevalent at all ages of the dentate population. Thus we aimed to investigate the association of salivary flow rate and pH with gingivitis and dental caries in patients of diabetes mellitus type II.

MATERIALS & METHODS:

This study is conducted in Madinah Medical Center Karachi. Subjects who are agreed to participate, filled the consent form and are all volunteers. The study received the ethical approval from Muhammad Bin Qasim Medical & Dental College Karachi. The inclusion criteria for the study is: age limit should be between 30-60 years, fasting blood glucose level should at least 140 mg/dl or more, presence of type II Diabetes for atleast 6 months, no other systemic disease other than diabetes.

Present study involved 100 adult human subjects after taking institutional ethical approval and informed consent. The subjects were divided into two groups.

Group I: control group contain 25 subjects having normal glucose level

Group II: Diabetic group (type I/ type II) contain 75

subjects. Fasting glucose level is measured by using glucometer.

Inclusion & exclusion criteria:

Patients with diabetes mellitus type II having age range was 30 -60 years. While patients having type II DM with other systemic diseases, taking any medicines, smokers etc are excluded.

Saliva collection:

Unstimulated saliva was collected for 5 minutes from participants using spitting method following preliminary mouth wash to water. Whole saliva was collected in sterile plastic bottle. The samples were transferred to Eppendorf tubes immediately after collection and stored at -20 °C until analysis.

Analytical method:

pH is immediately done before store the saliva in Eppendorf tubes . Electronic PH meter was used to measure the pH of saliva after calibration standards of 4 and 7. Samples with very low volume, pH was determined with pH paper strip. The flow rate was measured as mg/ml.

Analysis of Gingival Index and Dental Caries:

The criteria for gingival index was described according to the method of Suomi & Barbano 1968. Caries scores were classified according to ICDAS.

Statistical analysis:

Results were presented as mean $\pm$ SEM. To analyze the level of significance between flow rate and PH with gingivitis and dental caries ANOVA with post hoc test was used. $P < 0.05$ was considered to be significant. Pearson's correlation was used to detect the correlation between flow rate and PH with gingivitis and dental caries.

RESULTS:

Demographic analysis:

The demographic data showed that mean fasting glucose level is 263 ± 62.7 in diabetes subjects. Female subjects are 59.45 % while male subjects are 32 %. 46.42 % of male and 53.5 % of female diabetes subjects are claimed to gingivitis while 43.85 % of male and 56.0 % of female diabe-

Table I: Demographic analysis:

Parameters	Mean $\pm$ SD	Total %	Male%	Female%
Control	109 $\pm$ 11.9	25	11	16
Diabetes	263 $\pm$ 62.7	75.0	32	59.45
Salivary PH of diabetes	6.70 $\pm$ 0.96	75	32	59.45
Salivary flow rate of diabetes	0.307 $\pm$ 0.19	75	32	59.45
Gingival indices	2.28 $\pm$ 0.62	74.0	46.42	53.5
Dental caries indices	2.25 $\pm$ 0.61	72.7	43.85	56

Table II: Effect of Diabetes, salivary flow rate, salivary PH on Gingival and Dental caries indices:

PARAMETERS	GINGIVAL INDICES	DENTAL CARIES INDICES	P<0.05
control	0.00	0.00	significant
Diabetes	0.00,a n/s	0.00	significant
Salivary flow rate	0.00d/b n/s	0.00	significant
Salivary PH	0.00f/cn/s	0.00	significant

One way ANOVA showed significant results.

a,b,c represents: when compared gingival indices with dental caries; n/s : non-significant

tes subjects are claimed to dental caries as shown in table I.

Effect of Diabetes, salivary flow rate, salivary PH on Gingival and Dental caries indices:

We observed a significant results between salivary flow rate and pH with gingivitis. Similarly a significant differences were also observed between salivary flow rate and PH with dental caries as shown in table II.

Correlation between diabetes, salivary flow rate & salivary pH with gingivitis and dental caries:

Figure I showed a linear positive correlation of salivary flow rate with gingivitis and dental caries. A linear negative correlation is observed between salivary PH with gingivitis and dental caries. Strong and positive correlation is also observed between gingivitis and dental caries.

DISCUSSION

We found significant results of gingival and periodontal indices in diabetic patients. A significant results also found between salivary flow rate and PH of diabetes group. Diabetes is found to have several oral diseases including salivary and oral dysfunction, oral bacterial and fungal infections,




Figure I showed a linear positive correlation of salivary flow rate with gingivitis and dental caries.

oral mucosa lesions. We found significant association of diabetes with gingivitis and dental caries. Xerostomia is characteristic feature of diabetes mellitus patients, which may facilitate oral fungal infections.

Previously reports have emphasized the close relationship between diabetes and dental caries. We found significant result of diabetes and dental caries indices as shown in table II. We observed negative correlation between diabetes and dental caries (figure I). Nichols found the same result and they suggested that cariogenic diets containing large amounts of sugar or inoculation of infectious cariogenic bacteria into the oral cavity to induce aries in experimental animals. We found low salivary flow rate and PH of DM type II patients when compared to control as discussed in previously. The low saliva PH is strong evidence of reduced buffer salivary capacity and increased caries risk. The acidic PH of saliva leads metabolic acidosis which in turn loss the protective mechanism of saliva in diabetes patients. This acidic PH promotes the growth of aciduric bacteria and allows the acidogenic bacteria to proliferate creating an inhospitable environment for the protective oral bacteria. Oral environmental balance is changed and is favorable for cariogenic bacteria, lowers the PH further and the cycle is continues.

We found significant results in gingivitis indices when compared with positive control. One of the major complications of diabetes is change in microvascular integrity. Advanced glycation end products are formed irreversibly in chronic diabetes mellitus patients in the tissues and changes the collagen stature, altered immune function. This lead impairment of polymorphonuclear leukocyte function and facilitate bacterial proliferation in the tissues. Causes increased proinflammatory IL 1 β , TNF- α leads to increase collagenase activity and decreased collagen synthesis. This results in compromised wound healing and oral health.

A negative correlation is found between pH of saliva with dental caries and gingivitis indices which suggested that when pH of saliva decreased there is high chances of dental caries and gingivitis. While

we observed positive and strong correlation of gingivitis and dental caries. Therefore, it is highly likely that severe gingivitis or marginal periodontitis might have occurred due to severe periapical inflammation due to dental caries in male diabetic WBN/KobSlc rats.

CONCLUSION:

we found significant decreased level of salivary flow rate and pH in diabetes mellitus patients. A significant differences were observed when compared salivary flow rate with gingivitis and dental caries indices. We also observed significant differences of salivary PH with gingivitis and dental caries indices when compared. A negative correlation was observed between salivary pH with gingivitis and dental caries indices while, positive and strong correlation is observed between gingivitis and dental caries indices. It is therefore concluded, that diabetes mellitus is directly effect on oral health and gingivitis is secondary to dental caries.

REFERENCES

- 1 Yuen HK. Factors associated with additional time dental hygienists spent on educating patients with diabetes. Spec Care Dentist. 2018 Jul 6. doi: 10.1111/scd.12312
- 2 Grossi SG. Treatment of periodontal disease and control of diabetes: an assessment of the evidence and need for future research. Ann Periodontol. 2001; 6(1): 138-45.
- 3 Dental caries; the disease and its clinical management. Copenhagen, Denmark: Blackwell Munksgaard;2003.
- 4 Field EA, Longman LP, Buchnall R, Kaye SB, Higham SM, Edgar WM. The establishment of a xerostomia clinic: a prospective study. Br J Oral Maxillofac Surg. 1997; 35(2): 26-103.
- 5 Chavez EM, Borrell LN, Taylor GW, Ship JA. A longitudinal analysis of salivary flow in control subjects and older adults with type 2 diabetes. Oral Surg Oral Med Oral Pathol Oral Radiol Endod. 2001 Feb;91(2):166-73.
- 6 Dye BA, 2012. Global periodontal disease epidemiology. Periodontol. 2000(58): 10-25A
- 7 Al Bandar JM. Periodontal diseases in North America. Periodontol 2000. 2002; 29: 31-69.
- 8 Suomi, J. D. & Barbano, J. P.: Patterns of gingivitis, J Pedodontol, 39-71-74, 1968.
- 9 International Caries Detection and Assessment System Coordinating Committee. International Caries Detection and

Assessment System (ICDAS II): Criteria Manual. Revised in December and July 2009 Bogota, Colombia and Budapest, Hungary.

10 Saini R, Al-Maweri SA, Saini D, Ismail NM, Ismail AR. Oral mucosal lesions in non oral habit diabetic patients and association of diabetes mellitus with oral precancerous lesions. *Diabetes Res Clin Pract.* 2010;89:320–326.

11 Fernando Llambés, Santiago Arias-Herrera, and Raúl Caffesse. Relationship between diabetes and periodontal infection. *World J Diabetes.* 2015 Jul 10; 6(7): 927–935.

12 Borghelli RF, Devoto FC, Lazzari RN, Erausquin J, Foglia VG. 1971. Effect of insulin on caries activity in the rat. *J Dent Res* 50:888–891

13 Nichols MS, Shaw JH. 1957. The effect of alloxan diabetes on caries incidence in the albino rat. *J Dent Res* 36:68–74

14 Siudikiene J, Machiulskiene V, Nyvad B, Tenovou J, Nedzelskiene I. Dental caries and salivary status in children

with type 1 diabetes mellitus, related to the metabolic control of the disease. *Eur J Oral Sci* 2006; 114: 8-14.

15 Prathibha KM, et al. Evaluation of salivary profile among adult type 2 diabetes mellitus patients in South India. *J Clinical and Diagnostic research.* 2013; 7(8): 1592-95.

16 Goyal D, Kaur H, Jawanda MK, et al. Salivary pH and dental caries in diabetes mellitus. *International Journal of Oral & Maxillofacial Pathology.* 2012;3(4):13–16.

17 Al-Maskari AY, Al-Maskari MY, Al-Sudairy S. Oral Manifestations and complications of diabetes mellitus A review. *SQU Med J.* 2011;11(2):179–86

18 Yasushi Kodama,1 Masahiro Matsuura,3 Tomoya Sano,2 Yutaka, Nakahara,2 Kiyokazu Ozaki,2 Isao Narama,2 Tetsuro Matsuura2,* Diabetes Enhances Dental Caries and Apical Periodontitis in Caries-Susceptible WBN/KobSlc Rats. *Comp Med.* 2011 Feb; 61(1): 53–59.